

Osteogenetic-Pneumopedic™ protocol:
in accord with the cranio-caudal gradient of human development

- **A cranial examination by a cranial osteopath of a young baby soon after birth
If cranial asymmetry is diagnosed, cranial osteopathy may be provided as indicated;**
- **An oral examination by a pediatric dentist or general dentist of a young baby soon after birth. If ankylosis of the tongue is diagnosed, lingual frenectomy may be provided by an oral surgeon;**
- **An oral myofunctional examination by a myofunctional therapist, by the age of three years. If myofunctional habits, such as digit-sucking, lip-sucking etc are diagnosed, MFT may be provided;**
- **An orthodontic examination by an orthodontist or dentist, by the age of three to six years. If a malocclusion is diagnosed or developing, osteogenetic-orthodontic therapy may be provided;**
- **A TMJ examination by a neuromuscular dentist or general dentist, by the age of eight years. If TMD is diagnosed or developing, Phase I TMD therapy may be provided as indicated;**
- **A cervical examination by a chiropractor, osteopath etc, by the age of nine years.
If cervical lordosis or kyphosis is diagnosed, therapy may be provided as indicated;**
- **A sleep study examination by a Sleep specialist, pulmonologist, ENT etc., by age three to six years. If sleep-disordered breathing is diagnosed, osteogenetic-pneumopedic™ therapy may be provided.**